

# Phytochemical Properties and Antibacterial Activity of Leaf Extract of *Ocimum gratissimum* On *Salmonella* Species

## Abstract

**Aim:** *Ocimum gratissimum* is commonly used as food and health purposes. This study is aimed at evaluating the bioactive compounds and antibacterial activity of leaf extract of *O. gratissimum* against *Salmonella* species.

**Methodology:** The Phytochemical screening of *O. gratissimum* was conducted using standard methods. Screening for antibacterial activity of the leaf extracts against *Salmonella* species was determined using agar well diffusion method. An *in-vivo* toxicity study was carried out with albino rats.

**Results:** The phytochemical screening revealed the presence of saponins, tannins, cardiac glycoside, flavonoid, glycosides, alkaloid, volatile oils and steroids. A zone of inhibition of 14mm was recorded against the organisms using ethanolic extract with a concentration of 100mg/ml and the lowest was recorded against *Salmonella paratyphi* with the concentration of 25mg/ml of the ethanolic extract. Zone of inhibition of 9.00mm and 10.0mm was recorded against *S. typhi* and *S. paratyphi* on a concentration of 100mg/ml of the aqueous extract. A minimum inhibitory concentration of 100mg/ml and 25mg/ml of the aqueous and ethanolic extract of the leaf was recorded. After the toxicity test, no death was recorded after 2 (two) weeks.

**Conclusion:** The leaf extract of *O. gratissimum* shows promising potentials in the treatment of infectious diseases associated with *Salmonella typhi* and *Salmonella paratyphi*, due to its antimicrobial activity and low toxicity. However, further studies are needed to non-polar solvents to isolate other bioactive compounds as well as identify the active metabolites responsible for these activities.

**KEYWORDS:** *Ocimum gratissimum*, antibacterial activity, salmonella typhi, salmonella paratyphi, phytochemical, toxicity.

## INTRODUCTION

34 Medicinal plants are known to contain, in one or more of its organs, substances that can be used for  
35 therapeutic purposes or as precursors for the synthesis of useful drugs [1]. Many of such plants  
36 known to be used primitively to alleviate symptoms of illnesses have been screened to have  
37 medicinal importance, some of which include: *Vernonia amygdalina* (bitter leaf), *Ocimum*  
38 *gratissimum* (scent leaf), *Zingiber officinale* (ginger), *Azadirachta indica* (Dogonyaro), *Piper*  
39 *guineense* (lyere), *Allium sativum* (garlic), Cotton leaf (*Gossypium* spp) etc. These plants have been  
40 reportedly used in the traditional treatment of ailments such as stomach disorder, fever symptoms  
41 and cough [2].

42 Researchers are increasingly turning their attention to natural products looking for new leads to  
43 develop better drugs against cancer, as well as viral and microbial infections. The phytochemical  
44 evaluation of *Ocimum gratissimum* shows that it is rich in alkaloid, tannins, oxalate, flavonoids and  
45 essential oil [3]. In the coastal area of Nigeria, the plant *Ocimum gratissimum* is used in the  
46 treatment of epilepsy, high fever and diarrhoea [4]. *Ocimum gratissimum* (Scent leaf) is a perennial  
47 plant which is widely distributed in the tropics of Africa and Asia. It belongs to the family Labiatae  
48 and it is the most abundant of the genus *Ocimum*. In the southern part of Nigeria, it is called “Efirin  
49 nla” by the Yoruba speaking tribe. “Nichonwu” in Igbo while in the northern part of Nigeria, it is  
50 called “Daidoga” [5]. Leaf extract of *Ocimum gratissimum* and *Xylopiya aethiopiaea* were analyzed  
51 against five pathogenic organisms. *Staphylococcus aureus*, *Escherichia coli*, *Streptococcus*  
52 *fecalis*, *Pseudomonas aeruginosa* and *Lactobacilli* [3]. The findings justify the application of *Ocimum*  
53 *gratissimum* in dermatological cream and indicate the effective doses could be achieved at very low  
54 concentration and also shows that the aqueous fractions of both plants have more potential as  
55 antimicrobial agents than their ethanolic fractions [3]. The findings of Silva *et al.* [6] showed the  
56 extracts of *Ocimum gratissimum* to be active against human pathogenic dermatophytes.

57  
58 A thousand years ago an extensive use of plants as medicines has been reported and was initially  
59 taken in the form of crude drugs such as tinctures, elixirs, poultices, powders, and other herbal  
60 formulations [7]. However, the use of herbal products should be based on scientific origin; otherwise  
61 they would be useless and unsafe [7]. Furthermore, the irrational use of these herbal products may  
62 cause serious toxicity for humans. Unfortunately, many people underestimate the toxicity of natural  
63 products and do not realize that these agents could be as toxic as or more toxic than synthetic  
64 drugs [7]. A typical example of a toxic herbal product is the leaves of *Atropa Belladonna* and  
65 *Digitalis purpurea* [8], which show severe systemic toxicity if taken orally.

66  
67 Toxicology is the important aspect of pharmacology that deals with the adverse effect of bioactive  
68 substance on living organisms prior to the use as drug or chemical in clinical use [9], As per the  
69 OECD 2001 guidelines, in order to establish the safety and efficiency of a new drug, toxicological  
70 studies are very essential in animals like mice, rat, guinea pig, dog, rabbit, monkey etc under

71 various conditions of drug. Toxicological studies help to make decision whether a new drug should  
72 be adopted for clinical use or not. OECD does not allow the use of drug clinically without its clinical  
73 trial as well as toxicity studies. The aim of this present work, therefore, was to carry out  
74 phytochemical screening of the leaf of *Ocimum gratissimum*, study the antibacterial effects of the  
75 leaf extracts of *Ocimum gratissimum* on selected Enterobacteriaceae (*Salmonella* species) and to  
76 estimate the toxic effects of aqueous and ethanolic extracts from *Ocimum gratissimum* in albino  
77 Rats.

## 78 **MATERIALS AND METHODS**

### 79 **Collection and Identification of Leaf Materials**

80 *Ocimum gratissimum* (scent leaf) was obtained from Meat Market, Sokoto, Nigeria. The collected  
81 leaf was identified and authenticated at the Herbarium Section of the Department of Biological  
82 Sciences, Botany Unit of Usmanu Danfodiyo University Sokoto, Sokoto State, Nigeria. Voucher  
83 specimen numbers UDUH/ANS/101 was obtained.

### 84 85 **Preparation and Extraction of Leaf Extracts**

86 The fresh leaves were allowed to dry completely at room temperature before using them for this  
87 study. The leaf material was pulverized using mortar and pestle into a fine powder. Two different  
88 solvents were used for the extraction namely: water and ethanol. A 100g of powdered leaf was  
89 soaked in 1000ml of each solvent in accordance with Udochukwu *et al.* [10]. Each solution was  
90 stirred intermittently and allowed to stand for 48h, and then filtered by first, using a clean muslin  
91 cloth and then, No. 1 Whatman filter paper. Sterilization of the solutions was made using membrane  
92 filters. The sterile extract obtained was stored in sterile capped bottles and refrigerated [11].

### 94 **Characterization and Identification of *Salmonella* Species**

#### 95 **Source of Test Organism**

96 The test organisms for this study (*Salmonella* species) are members of the family  
97 Enterobacteriaceae. The pure clinical isolates of *Samonella typhi* and *Samonella paratyphi* were  
98 obtained from the Department of Medical Microbiology and Parasitology, Specialist Hospital Sokoto,  
99 Nigeria. All the clinical isolates were checked for purity by sub-culturing the isolates onto  
100 Salmonella-Shigella Agar medium. After 24hrs of incubation, there were growths of the isolates and  
101 they were maintained on nutrient agar slants at 4<sup>0</sup>C in the refrigerator until required for further use.

#### 104 **Biochemical Confirmation and Serotyping of *Salmonella***

106 The ISO-6579 [12], the standard recommendation was used for biochemical confirmation of  
107 *Salmonella*. The subculture of characteristic colonies from each Petri dish of Salmonella-Shigella  
108 agar medium was made. The triple sugar iron agar (TSI agar), Urea agar/broth, L-lysine

109 decarboxylase,  $\beta$ -galactosidase (ONPG), Voges Proskauer and Indole tests were followed in this  
110 order.

111  
112 In serotyping, the subculture of characteristic colonies from each Petri dish of Salmonella-Shigella  
113 agar was transferred onto nutrient agar slopes and incubated overnight at 37°C. Using a wire loop,  
114 3 separate drops (each 0.02 ml) of saline solution were placed onto a clean microscope slide.  
115 Growth from the agar slope was added and emulsified to produce homogeneous suspension. A  
116 loopful of *Salmonella* polyvalent 'O' (PSO) anti-serum was mixed with the first drop of suspension  
117 and a loopful of *Salmonella* polyvalent 'H' (PSH) anti-serum with the second drop. It was rocked  
118 gently back and forth and examined for agglutination against a black background. Positive results  
119 were recorded if agglutination occurred within 20 min after shaking against dark background. In  
120 order to exclude any spontaneous agglutination (auto-agglutination), a negative control (using  
121 physiological saline solution and bacterial colony to be tested) was included in the test.

### 122 123 **Standardization of Bacteria Cell Suspension**

124 The nutrient broth cultures of the organisms for this study were taken and inoculated at 37°C on a  
125 fresh agar plate of nutrient agar for 24 hours. Sterile distilled water (2ml) was poured on it and then  
126 mixed with the inoculums, 1ml of each was taken and transferred into 9ml of sterile distilled water  
127 and diluted to 0.5 Macfarland Standard giving a load of  $10^5$ -  $10^6$  organisms/ml. One hundred  
128 microlitres of these were taken and poured onto the surface of the agar and then spread evenly with  
129 the use of a spreader on the plate to be used for the study [13].

### 130 131 **Preparation of Extracts Concentration**

132 The different extracts of the sample were reconstituted with sterile distilled water. The initial  
133 concentration of each plant extracts (1g) was diluted using 10ml of sterile water to obtain the stock  
134 culture. From this stock culture, different concentrations were gotten which were 100mg/ml,  
135 50mg/ml, 25mg/ml, 12.5mg/ml, 6.25mg/ml, and 3.125mg/ml for each of the extracts (water and  
136 ethanol).

### 137 138 **Determination of Antibacterial Activities of Leaf Extracts**

139  
140 Agar-well diffusion Method was employed for the antibacterial testing [14]. The antibacterial  
141 screening of the extracts was done as described by Perex *et al.* [14]. One (1) gram of each crude  
142 extract (aqueous and ethanolic) was poured into 10ml water. From this stock culture, different  
143 concentrations were gotten which were 100mg/ml, 50mg/ml, 25mg/ml, 12.5mg/ml, 6.25mg/ml, and  
144 3.125mg/ml for each of the extracts (aqueous and ethanol). Nutrient agar was poured in sterile Petri  
145 dishes and was allowed to solidify. A loopful of the test culture of MacFarland standard was dropped  
146 on the solidified agar and the organism was spread all over the surface of the agar using a spreader  
147 (wire loop). The inoculated plates were allowed to dry after which wells of approximately 5mm in  
148 diameter were made on the surface of the agar medium using a sterile cork borer. Then, 0.2ml of

149 different concentrations of the extract was separately introduced into the different wells that have  
150 been labelled accordingly. This procedure was repeated in triplicate and allowed to stay for 30mins  
151 on the bench after which they were incubated for 24h at 37°C. At the end of incubation, observed  
152 zones of inhibition were measured and recorded to the nearest millimetre.

### 153 154 **Determination of Minimum Inhibitory Concentration of the Extracts** 155

156 This was carried out using the agar diffusion method following the recommendations of the Clinical  
157 and Laboratory Standard Institute [15]. Different concentrations 100, 50, 25, 12.5, 6.25 and 3.125  
158 mg/ml of the extracts were prepared and 1ml from each of the concentrations of the extracts was  
159 added onto molten nutrient agar and was mixed thoroughly. Then, 1µl of an overnight nutrient broth  
160 culture of the test isolates were added to each plate of the Mueller-Hinton agar containing the  
161 extracts and incubated at 37°C for 24 h. The experiment was conducted in triplicate for all the test  
162 isolate. Plates without visible growth of the organisms in each concentration were taken as the MIC  
163 [11].  
164

### 165 **Phytochemical Screening of Leaf of *Ocimum gratissimum***

166 The pulverized leaf obtained was subjected to phytochemical screening to determine the presence  
167 of bioactive compounds.  
168

#### 169 **Test for Tannins**

170 Five per cent (5%) ferric chloride was added drop by drop to 3ml of each extract and observed for  
171 brownish green or a blue-black colouration [16].  
172

#### 173 **Test for Saponins**

174 Two grams (2g) of the powdered sample of each extract was boiled in 20ml of distilled water in a  
175 water bath and filtered. Then, 10ml of the filtrate was mixed with 5ml of distilled water and shaken  
176 vigorously for a stable persistent froth. The frothing was mixed with 3 drops of olive oil, shaken  
177 vigorously and then observed for the formation of emulsion [17].  
178

#### 179 **Test for Flavonoids**

180 One millilitre (1ml) of 10% NaOH solution was added to a portion of the aqueous filtrate of each  
181 plant extract, followed by addition of concentrated H<sub>2</sub>SO<sub>4</sub>. A yellow colouration observed in the  
182 extract indicated the presence of flavonoids [16].  
183

#### 184 **Test for Cardiac Glycosides**

185 Five millilitres (5ml) of each extract was treated with 2ml of glacial acetic acid containing 1 drop of  
186 ferric chloride solution (3.5%). The content was allowed to stand for one minute. One millilitre (1ml)

187 of concentrated H<sub>2</sub>SO<sub>4</sub> was carefully poured down the wall of the tube. A reddish-brown ring of the  
188 interface indicated a deoxysugar characteristic of cardenolides [18].

189  
190 **Test for Alkaloid**

191 Two milliliter (2ml) of each extract was stirred with 2ml of 10% dilute hydrochloric acid. Then, 1ml  
192 was treated with a few drops of Wagner's reagent and second 1ml portion treated with Mayer's  
193 reagent. Deep brown precipitation indicated a positive test [18].

194  
195 **Test for Glycosides**

196 The 2.5ml of 50% H<sub>2</sub>SO<sub>4</sub> was added to 5ml of each of the extracts in test tubes. The mixture was  
197 heated in boiling water for 15minutes. Cooled and neutralized with 10% NaOH, 5ml of Fehling's  
198 solution was added and the mixture was boiled again. A brick-red precipitate was observed, which  
199 indicated the presence of glycosides [18].

200  
201 **Test for Steroids**

202 This was carried out according to the method of Harborne [18]. One (1) ml of each leaf extract was  
203 added in 2ml of chloroform, and 2ml of sulphuric acid (H<sub>2</sub>SO<sub>4</sub>) was added thereafter. A red  
204 colouration confirmed the presence of steroids.

205  
206 **Test for Volatile oils**

207 One millilitre (1ml) of each of the extract fractions was mixed with 5ml of dilute HCL. A white  
208 precipitate was formed, which indicated the presence of volatile oils [17].

209  
210 **Toxicity Study of the Leaf Extracts of *Ocimum gratissimum***

211  
212 Acute oral toxicity test was carried out using the procedure of the Organization for Economic  
213 Cooperation and Development [19]. Ten (10) randomly selected Albino rats were used. The rats of  
214 both sexes weighing 160-200g were used for the study. The animals were obtained from the Faculty  
215 of Veterinary Medicine, Usmanu Danfodiyo University, Sokoto. The animals were acclimatized for a  
216 period of seven days. All animals were housed, caged and allowed free access to food and water  
217 before they were used for the experiment. The animals' weights were taken and starved of food.  
218 Then 5000mg/kg bodyweight of the extract was administered in a single concentration.  
219 Concentrations were calculated according to the bodyweight of the animals. Oral administration of  
220 extracts was done using a graduated syringe and cannula. They were placed under observation for  
221 48 hours for behavioural changes and daily for 14days for mortality [19], upon which the number of  
222 deaths and LD<sub>50</sub> were determined.

223  
224 **RESULTS AND DISCUSSION**

225

226 The results of phytochemical screening of *O. gratissimum* leaves revealed the presence of the  
 227 following secondary metabolites; tannins, saponins, flavonoid, steroid, cardiac glycoside,  
 228 glycosides, alkaloid, and volatile oil. This is similar to the findings of Nweze *et al.* [20] who reported  
 229 the presence of alkaloids, tannins, glycoside, saponin, cardiac glycoside, steroid and flavonoids in  
 230 *O. gratissimum* which is similar to the results obtained in this study.

231 The results revealed that the aqueous extract of *Ocimum gratissimum* had less inhibitory activity on  
 232 the test organisms (Table 2), while the ethanolic extracts of *Ocimum gratissimum* (Table 3) had  
 233 antibacterial activity against the isolates tested. At 100mg/ml concentration, the ethanolic extracts  
 234 showed greater antibacterial activity than the aqueous extracts as indicated by zones of inhibition.  
 235 At 12.5mg/ml – 3.125mg/ml, the ethanolic extracts of *Ocimum gratissimum* (Table 3) was not  
 236 effective on the isolates. While at 50mg/ml – 3.125mg/ml the aqueous extracts of *Ocimum*  
 237 *gratissimum* (Table 2) was not effective on the isolates. This indicates that the antibacterial activity  
 238 of this leaf extracts is concentration-dependent. Ethanolic extract showed high inhibitory zones than  
 239 aqueous extracts and when compared to standard antibiotic such as Pemaclav drug had an  
 240 appreciable zone of inhibition of the test organisms. The result of this work showed that the  
 241 ethanolic extract showed high inhibitory zones than aqueous extracts. This observed difference  
 242 between these plants extracts may be due to insolubility of active compounds in water or the  
 243 presence of inhibitors to the antimicrobial components Okigbo and Ogbonnanya [21], Amadioha and  
 244 Obi [1999], Okigbo and Ajale [2005]. They have attributed this observation to the high volatility of  
 245 ethanol which tends to extract more active compound from the sample than water, hence, this study  
 246 follows similar trends. The aqueous extract of *O. gratissimum* showed a decrease in the level of  
 247 inhibition against isolates at the highest concentration compared to the positive control, inhibition  
 248 zones ranging from 9.0 to 10.0 mm.

249  
 250  
 251  
 252

**Table 1. Phytochemical constituents of the leaf of *O. gratissimum* Leaf Extract.**

Phytochemical	<i>O. gratissimum</i>
Tannins	+
Saponins	+
Flavonoid	+
Cardiac glycoside	+
Alkaloid	+
Glycosides	+

Steroid	+
Volatile oil	+

253  
254  
255  
256  
257  
258

**KEY:-** = Not detected, + = Detected

**Table 2. The Antibacterial Activities of Aqueous Leaf Extracts of *O. gratissimum***

Test Organism	Zone of inhibition(mm) <i>O. gratissimum</i>							
	Concentration (mg/ml)	100	50	25	12.5	6.25	3.125	+ve Control
<i>Salmonella typhi</i>	09.0	x	x	x	x	x	x	20.0
<i>Salmonella paratyphi</i>	10.0	x	x	x	x	x	x	20.0

259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269

**Key:**

Values are mean of three replicates (n=3)

x = No zone of inhibition

+ve control = Pemaclav drug (10mg/ml)

**Table 3. The Antibacterial Activities of Ethanolic Leaf Extracts of *O. gratissimum***

Test Organism	Zone of inhibition(mm) <i>O. gratissimum</i>							
	Concentration (mg/ml)	100	50	25	12.5	6.25	3.125	+ve control
<i>Salmonella typhi</i>	14.0	13.0	12.0	x	x	x	x	20.0
<i>Salmonella paratyphi</i>	14.0	11.0	07.0	x	x	x	x	20.0

270  
271  
272  
273

**Key:**

Values are mean of three replicates (n=3)

x = No zone of inhibition

+ve control = Pemaclav drug (10mg/ml)

274  
275  
276  
277  
278  
279  
280  
281  
282

The minimum inhibitory concentrations (MIC) of aqueous and ethanolic leaf extracts on the test organisms ranged between 25mg/ml –100mg/ml. The minimum inhibitory concentrations of ethanolic extracts of *O. gratissimum* as 25mg/ml while aqueous leaf extracts of *O. gratissimum* had their MIC as 100mg/ml. Minimum inhibitory concentrations (MICs) of both aqueous and ethanolic extracts on test organisms using agar dilution method revealed low MIC, which is an indication of high efficacy of the leaf extracts while high MIC may indicate low efficacy or possible development of resistance by the microorganisms to the antimicrobial [24]. The aqueous extract showed its MIC at high concentration of 100mg/ml while ethanolic extract showed its MIC at 25mg/ml.



283 Oral administration of a single dose of ethanol and aqueous extracts of *O. gratissimum* of  
 284 5000mg/kg bodyweight of the test animals produced no mortality in them. The general signs and  
 285 symptoms of toxicity were observed for a period of 14 days after administration of the extracts.  
 286 However, the following observations were made during the exposure period; slow movement,  
 287 scratching of hair and mouth, tremor, raised hair coat and weakness. Thus, the median dose (LD<sub>50</sub>)  
 288 of the leaf extracts was estimated to be greater than 5000mg/kg because 5000mg/kg is the highest  
 289 dose according OECD [19]. From the experiment performed as per the OECD Guidelines 2001, the  
 290 results reveal that both aqueous and ethanolic extract of *Ocimum gratissimum* has been found  
 291 nontoxic at 5000 mg/kg body weight of experimental animals as in the first 1 hour of observation, no  
 292 morbidity was observed but weakness, slow movement, scratching of mouth, fur and body, tremor  
 293 was observed and in the next 48 hours of observation mortality were not found and all that  
 294 parameters used for evaluation of toxicity were found to be normal. No significant changes were  
 295 observed in body weight. In the last 2 weeks of observation, no death rate recorded. As per  
 296 observations and calculations from Acute Oral Toxicity (OECD Guidelines 2001), the LD<sub>50</sub> value of  
 297 aqueous and ethanolic Extract of *Ocimum gratissimum* was found to be more than 5000 mg/kg body  
 298 weight of the rats.

299  
 300 **Table 4 Minimum Inhibitory Concentration of the Aqueous and Ethanol Leaf Extracts of *O.***  
 301 ***gratissimum* against *Salmonella* spp**  
 302

<b>Bacterial Isolates</b>	<b>Aqueous extract MIC (mg/ml)</b>	<b>Ethanol extract MIC (mg/ml)</b>	<b>Pemaclav drug (Amoxicillin combination) MIC (mg/ml)</b>
<b><i>Salmonella typhi</i></b>	100	25	12
<b><i>Salmonella paratyphi</i></b>	100	25	12

303 Values are mean of three replicates (n=3)  
 304  
 305  
 306  
 307

308 **Table 5. Acute Toxicity Results on Twenty Randomly Selected Albino Rats.**

<b>Dose (mg/kg)</b>	<b>Time Duration</b>	<b>No. of Animals</b>	<b>No. of Deaths</b>	<b>Observation Signs</b>
5000	0-30 minutes 1 hour 24 hours 48 hours	5	0	Weakness, slow movement immediately after administration. Continuously scratching of mouthpart, fur and body, tremor. Ruffled fur, scratching of their nostril. Normal movement and less scratching of the body

	2 weeks			part. No death rate recorded.
5000	0-30 minutes 1 hour 24 hours 48 hours 2 weeks	5	0	Increased breathing  Scratching of mouth and body parts Ruffled fur No scratching of the body part No death rate recorded

309 Key:

310 a) The first 5 rats were given aqueous leaf extracts of *O. gratissimum*

311 b) The last two 5 rats were given ethanolic leaf extracts of *O. gratissimum*

312

### 313 Conclusion

314

315 From this study, it was observed that ethanol extracts exhibited high inhibitory activity on the test  
316 organisms. This can be deduced to the ability of ethanol to extract more of the essential oils and  
317 secondary plant metabolites which are believed to exert antibacterial activity on the test organisms.  
318 This suggests the possibility of using the ethanol extracts of *O. gratissimum* in treating the diseases  
319 caused by the test organisms. Aqueous and ethanolic extracts of *Ocimum gratissimum* exhibit no  
320 toxic effects when given orally at concentration of 5000mg/kg body weight. However, the normalcy  
321 and insignificant changes in toxicity parameters and body weights reveal the safety of aqueous and  
322 ethanolic extract at a dose of 5000 mg/kg body weight. This study, however, can justify the use of  
323 the leaf in traditional medicine practice as a therapeutic agent and can explain the long historical  
324 use of these plants.

### 325 REFERENCES

326

327 1. Sofowora A. *Medicinal plant and traditional medicine in Africa*. 3rd edition spectrum Books  
328 Limited, Ibadan, 1999; pp.172-188.

329 2. Odugbemi T. *Outline of Medicinal Plants in Nigeria*. 1st Edition, University of Lagos Press,  
330 Nigeria, p.77. 2006.

331 3. Ijeh II, Njoku OU Ekenza EC. Medicinal Evaluation of *Xylopi aethiopica* and *Ocimum*  
332 *gratissimum*. *Journal of Medicial Aromatis Science*. 2004; 26(1): 44-4

333

334 4. Sofowora A (1993). *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Books Ltd., Ibadan,  
335 Nigeria, pp. 191-289.

336

337 5. Effraim KD, Jacks TW, Sapiro OA. Histopathological Studies On the Toxicity of  
338 *Ocimum gratissimum* leaves extracts on some organs of Rabbits: *African Journal of*  
339 *Biomedical Research*. 2003; 6: 21-25.

- 340 6. Silva MRR Oliveira Fernandes OFL Passos XS. Instituto de Patologia Tropicale Saude  
341 Publica DaUFG, Goiania, Goias, Brazil. 2005.
- 342 7. Gullo VP, McAlpine JKin LS, Baker D Petersen F. An acute oral toxicity study of methanolic  
343 extract from *Tridax procumbens*. *Journal of Industrial Microbiology and Biotechnology*. 2006;  
344 33: 523-531.
- 345 8. Tripathi KD. *Essentials of Medical Pharmacology*, Jaypee Brothers Medical Publishers (P)  
346 Ltd., New Delhi, 6th edition. 2008; 455-465
- 347 9. Aneela S, Somnath DE, Lakshmi KK, Choudhury NSK, Das SL Sagar KV. International  
348 Journal of Research in Pharmacy and Chemistry. 2011; **1(4)**: 820-824.
- 349 10. Udochukwu U Omeje FI Uloma IS Oseiwe FD. Phytochemical analysis of *Vernonia*  
350 *amygdalina* and *Ocimum gratissimum* extracts and their antibacterial activity on some drug-  
351 resistant bacteria. *American Journal of Research Communication*. 2015; **3(5)**: 225-235.
- 352 11. Ronald MA. Microorganisms in our World. Mosby Year Book, Inc. St. Louis. 1995; p.765.
- 353 12. ISO 6579: Microbiology 4<sup>th</sup> ed. - General guidance on methods for the detection of  
354 *Salmonella*, International Organization for Standardization, Geneve, Switzerland. **6**: 66-78.  
355 2002.
- 356 13. Farrouq AA, Muomora GD, Magashi AM, Jodi SM, Nata'ala MK and Habibu KB.  
357 Phytochemical, Antibacterial and Toxicity Study of Leaf Extracts of *Vernonia amygdalina*,  
358 Delile on *Salmonella* Species. *Asian Journal of Research in Medical and Pharmaceutical*  
359 *Sciences*. 2019; 6(4): 1-8.
- 360 14. Perex CC Paul ML Bazerque P. Antibiotic assay by agar well diffusion method. *Experimental*  
361 *Medicine and Biology*. 1990; **15**: 113-115.
- 362 15. Clinical and Laboratory Standard Institute (2006). Methods for antimicrobial susceptibility  
363 tests for bacteria that grow aerobically. Approved Standard, Seventh Edition. Document M7-  
364 A7 (replaces M7-A4) Villanova PA. 3-11; 2006.
- 365 16. Harbone JB. Phytochemical methods. 1st Edition Chapman Halls, London, pp. 20–22. 1984.
- 366 17. Trease GE Evans WC. *Pharmacognosy*. 14<sup>th</sup> edn. Brailier Titidel Canada Macmillan  
367 publisher. 1989; p:530
- 368 18. Harborne JB. Phytochemical methods. A guide to modern techniques of plant analysis 2<sup>nd</sup>  
369 edition. Chapman and Hall London 1-13. 1973.
- 370 19. Organization for Economic Cooperation and Development: Guideline for testing chemicals.  
371 Acute Oral Toxicity- Acute Toxicity Class Method 423. Adopted 17th December, 2001. 1-26.  
372 2001.
- 373 20. Nweze EI Okafor JI Njoku O. Antimicrobial activities of Methanolic Extracts of *Trema*  
374 *guineensis* (Schumm and Thorn) and *Morinda lucida* Benth used in Nigerian Herbal  
375 *Medicinal Practice*. *J. Biol. Res. Biotechnol*. 2004; 2(1):39-46.

- 376 21. Okigbo RN and Ogbonnanya OU. Antifungal Effects of Two Tropical Plants Extracts *Ocimum*  
377 *gratissimum* and *Afromaomum melegueta* on Post-Harvest Yam *Discorea* spp rot. African  
378 Journal of Biotechnology. 2006; 5(9): 727-731.
- 379 22. Amadioha AC and Obi VI. Control of anthracnose disease of cowpea by *Cymbopogon*  
380 *citrates* and *Ocimum gratissimum*. Acta Phytopathologica Entomologica Hungarica. 1999; 34  
381 (1-2): 85-89.
- 382 23. Okigbo RN Ajale AN. Inhibition of some human pathogens with the tropical plant extracts  
383 *Chromolineena odorata* and *Citrus aurantifolia* and some antibiotics. *International Journal of*  
384 *Molecular Medicine and Advance Science*. 2005; 1:34-40
- 385 24. Shanmugam AT Bhuvanewari RA Nazeer S Sambasivam and S Vairamani *et al.*  
386 Pharmacological properties of the venom of a marine gastropod *Babylonia spirata* (L.).  
387 Journal of Pharmacology and Toxicology. 2008; 3: 222-229.  
388